



1 Tracking of injected and resident (previously injected) bacterial  
2 cells in groundwater using ferrographic capture

3 William P. Johnson\*, William O. McIntosh

4 *Department of Geology and Geophysics, University of Utah, Salt Lake City, UT 84112, USA*

5 Received 20 September 2002; received in revised form 7 December 2002; accepted 4 January 2003

7 **Abstract**

8 A high-resolution bacterial tracking technique, ferrographic capture, was used to enumerate fluorescent-stained bacterial  
9 cells that were injected into groundwater during a field experiment. The goal of the experiment was to investigate whether  
10 detachment of previously injected stained resident cells attached to aquifer sediment was enhanced in the presence of the newly  
11 injected mobile cells. This injection was an improvement on past experiments in that the attached (resident) cells were stained,  
12 allowing their concentrations to be enumerated directly by ferrographic capture (upon detachment). Contrary to expectations  
13 based on previous experiments, enhanced detachment of stained resident cells did not occur upon the arrival of injected cells.  
14 Consistent with previous experiments, however, was the observation of ephemeral increases in unstained cell concentrations  
15 coincident with the arrival of the stained injected cells. The ephemeral pulses of unstained cells were previously speculated to  
16 represent enhanced detachment of unstained indigenous cells in response to hydrodynamic collision with injected cells. The  
17 lack of enhanced detachment of stained resident cells in the present experiments indicates that increased concentrations of  
18 unstained cells may have occurred by mechanisms other than hydrodynamic collision. Visually observed variations in stain  
19 intensity indicated that increased unstained cell concentrations may have resulted from cell division at the low-concentration  
20 fringe of the injected plume.

21 © 2003 Elsevier Science B.V. All rights reserved.

23 *Keywords:* Ferrographic capture; Bacterial cells; Hydrodynamic collision

24 **1. Introduction**

27 Bacterial attachment and detachment during trans-  
28 port in porous media is a subject of interest to those  
29 concerned with filtration for water treatment (Harvey  
30 and Garabedian, 1991), pathogen transport in ground-  
31 water (Harvey, 1997), and in situ bioaugmentation  
32 (Steffan et al., 1999). Bacteria are colloidal, being in

the micrometer size range, and despite their greater  
33 complexity, their transport is governed by many of the  
34 mechanisms that govern the transport of mineral  
35 colloids. Mechanisms controlling colloid detachment  
36 have been less studied relative to those governing  
37 attachment, likely because rates of colloidal detach-  
38 ment tend to be orders of magnitude lower than rates  
39 of attachment. The large difference in rates of attach-  
40 ment versus detachment allows the exclusion of  
41 detachment in models describing the gross transport  
42 behavior of colloids (e.g. filtration theory). However,  
43 second-order aspects of colloid transport are con-  
44

\* Corresponding author.

*E-mail address:* [wjohnson@mines.utah.edu](mailto:wjohnson@mines.utah.edu) (W.P. Johnson).

45 trolled or influenced by detachment, e.g. bacterial  
46 tailing following an input pulse (Johnson et al.,  
47 1995; Zhang et al., 2001), and the profile of attached  
48 cell concentrations versus distance (Zhang et al.,  
49 2001). Among the many processes potentially influ-  
50 encing colloidal detachment is hydrodynamic colli-  
51 sion between mobile and attached colloids, which has  
52 been observed for microspheres in impinging jet flow  
53 experiments (Dabros, 1989; Dabros and van de Ven,  
54 1992) and bacteria and microspheres in parallel plate  
55 chambers (Meinders et al., 1995).

56 That hydrodynamic collision between mobile and  
57 attached particles can enhance detachment of attached  
58 particles has been established based on force balance  
59 calculations (Dabros, 1989; Dabros and van de Ven,  
60 1992), which indicate that hydrodynamic interaction  
61 between deposited and flowing particles in stagnation  
62 point flow systems causes significant variations of the  
63 normal force acting on the deposited particles even for  
64 relatively large particle separations, resulting in the  
65 escape of weakly bound particles. Bacteria are rela-  
66 tively “soft” particles, which thereby decreases their  
67 potential for hydrodynamic collision relative to  
68 “hard” mineral colloids. However, after observing  
69 increased detachment of attached microspheres and  
70 attached bacteria in the presence relative to the  
71 absence of mobile bacteria or microspheres, Meinders  
72 et al. (1995) concluded that hydrodynamic collision  
73 operated in their experiments in impinging jet flow  
74 systems.

75 Tentative observation of enhanced detachment of  
76 attached cells in the presence of mobile cells was  
77 recently reported from a bacterial transport study  
78 conducted in Oyster, VA (Johnson et al., 2001). In  
79 this past study (conducted July 2000), two strains of  
80 bacteria were injected that had been previously iso-  
81 lated from the subsurface of the Oyster site (DeFlaun  
82 et al., 1990). The isolates were stained with a vital  
83 fluorescent stain (Fuller et al., 2000) and were injected  
84 into established flow cells at the site. Injected cells  
85 were distinguished from resident cells (unstained  
86 indigenous cells) on the basis of their internal stain.  
87 Ephemeral increases in concentrations of unstained  
88 bacteria coincident with arrival of the stained bacteria  
89 were observed in several wells during the transport  
90 experiment. The unstained cells could have originated  
91 from either the injected cell population or the pop-  
92 ulation indigenous to the aquifer (not injected). To

93 originate from the injected cell population, the  
94 unstained cells would need to have been selectively  
95 concentrated relative to stained cells during transport.  
96 This could have occurred by division of injected cells  
97 (assuming lack of stain transfer to daughter cells), loss  
98 of stain by diffusion during transport, and lesser  
99 adhesion of unstained relative to stained cells. Cell  
100 division and stain loss in samples collected during  
101 peak breakthrough were insufficient to explain the  
102 observed pulses of unstained cells. Standard adhesion  
103 assays indicated no difference in adhesion of stained  
104 versus unstained cells. Furthermore, to explain the  
105 observed ephemeral dominance of unstained cells,  
106 selective concentration would need to have occurred  
107 exclusively on the low-concentration fringes of the  
108 bacterial plume. Based on the above observation, it  
109 was tentatively concluded that the unstained cells  
110 originated from the cell population indigenous to the  
111 aquifer (not injected). Potential mechanisms of  
112 appearance of unstained indigenous cells include  
113 growth or detachment in response to the arrival of  
114 injected cells, with detachment more likely, given the  
115 lack of a lag time between arrival of the unstained and  
116 injected cells.

117 A significant drawback of the previous study was  
118 reliance on indirect means to quantify unstained cells,  
119 i.e. unstained cells were quantified by the difference  
120 between stained and total cells (stained plus un-  
121 stained). In the present study, potential enhanced  
122 detachment of cells from the site sediment, in  
123 response to the arrival of mobile injected cells, was  
124 monitored directly since the attached cells were  
125 stained. It was determined in July 2001 that previ-  
126 ously injected cells (remaining from the July 2000  
127 injection) remained intact and visibly stained in the  
128 site groundwater, and it was assumed that these cells  
129 also resided on the sediment. In July 2001, the same  
130 two bacterial strains were injected as in the previous  
131 year, however, the cells were stained oppositely from  
132 the cells injected in the previous year (the July 2000  
133 injection). Switching the stains allowed for direct  
134 observation of the stained cells remaining from the  
135 July 2000 injection (hereafter referred to as stained  
136 resident cells).

137 Detachment of attached cells by hydrodynamic  
138 collision with mobile cells is thought to require  
139 generation of sufficient normal force to remove the  
140 attached cells. The magnitude of the normal force

141 resulting from hydrodynamic collision is directly  
142 proportional to the rate of shear due to flow in the  
143 system (Dabros and van de Ven, 1992). In experi-  
144 ments described by Dabros (1995) and Dabros and  
145 van de Ven (1992), the wall shear rates were in the  
146 range of 100–1000 s<sup>-1</sup>. In the experiments per-  
147 formed by Meinders et al. (1995), which examined  
148 rates of microsphere and bacteria detachment, the  
149 shear rates were ~ 50 s<sup>-1</sup>. The shear rates in our  
150 experimental systems are calculated to be about 5–10  
151 s<sup>-1</sup>, based on idealized parabolic pore geometry  
152 (Bergendahl and Grasso, 2000) and a representative  
153 grain size of 300 µm. These calculated values are 5–  
154 10 times less than those examined by the above-  
155 described studies (Dabros, 1995; Dabros and van de  
156 Ven, 1992; Meinders et al., 1995). However, in our  
157 system of packed porous media, the calculated values  
158 of shear likely underestimate the actual values for  
159 several reasons: (1) the grain size is distributed,  
160 thereby decreasing the pore throat sizes below those  
161 expected for homogenous media that comprised of  
162 300-µm grains; (2) the grains are not smooth nor  
163 spherical, and thereby do not result in the parabolic  
164 pore geometry idealized for our calculation.

165 Because the hypothesized collisions are hydrody-  
166 namic rather than direct, it may not be appropriate to  
167 assess the potential rate of hydrodynamic collision  
168 based on well-known rates of direct collision based on  
169 filtration theory (e.g. Ryan and Elimelech, 1996).  
170 However, given the above caveat, the rates of direct  
171 collision (per meter of transport) based on filtration  
172 theory were 60 and 850 for assumed average grain  
173 sizes of 500 and 100 µm, respectively. A range is  
174 given since the grain size of the natural sediment is  
175 distributed (the sediment displays an arithmetic mean  
176 grain size of 270 µm). The ranges in collision number  
177 obviously do not account for the packing effects of  
178 distributed grain sizes, but give some idea of the  
179 potential frequency of bacterial collision with sedi-  
180 ment grain surfaces. The unstained cell concentrations  
181 in the pulses observed in the previous experiment  
182 typically ranged from several hundred to several  
183 thousand cells per milliliter, which appears to be  
184 supportable via collision numbers between 60 and  
185 850 (per mobile cell). However, it must be stressed  
186 that the utility of the collision number in assessing the  
187 feasibility of hydrodynamic collision is limited, since  
188 the “collisions” that are hypothesized to occur bet-

189 ween mobile and attached bacteria are hydrodynamic  
190 rather than direct.

191 There is no simple means known to the authors of  
192 accurately determining the values of shear or the  
193 potential frequency of hydrodynamic collision in a  
194 packed porous media. Hence, the opportunity of a  
195 direct test for enhanced detachment of resident cells  
196 was useful. If enhanced detachment of resident stained  
197 bacteria had been observed, then further investigation  
198 of its causes would have been warranted. In contrast,  
199 the observed lack of enhanced detachment reported  
200 below indicates the need to further consider other  
201 mechanisms of appearance of the unstained cells.  
202 Regardless of outcome, the investigation described  
203 below demonstrates the utility of the high-resolution  
204 technique that was used for tracking the bacteria.

## 2. Methods

205  
206 Two bacterial strains were originally isolated from  
207 the site and were used in this study. DA001 is an  
208 aerobic adhesion-deficient variant selected using col-  
209 umn assays, and is identified as a *Comamonas* sp.  
210 (DeFlaun et al., 1990). DA001 is Gram-negative and  
211 is 1.2 × 0.6 µm in size. OY-107 is a facultative iron-  
212 reducing bacterium of the genus *Acidovorax* that was  
213 determined to be naturally adhesion deficient upon  
214 isolation from the South Oyster (SO) site. OY-107 is a  
215 Gram-negative bacterium of size 1.9 × 1.0 µm.

216 Bacterial strains were grown by Envirogen (Law-  
217 renceville, NJ) in minimal media supplemented with  
218 lactate using standard fermentation procedures.  
219 Strains were harvested by centrifugation and starved  
220 in at a cell concentration of approximately 10<sup>9</sup> cells/  
221 ml in artificial site groundwater for at least 48 h  
222 (DeFlaun et al., 2001). Cells were stained during  
223 preparation by Envirogen using vital fluorescent  
224 stains. Vital fluorescent stains cause no adverse effect  
225 to cell culturability (Fuller et al., 2000; Fuller et al.,  
226 2001a,b). DA001 was stained using the green fluo-  
227 rescent stain 5-(and 6)-carboxyfluorescein diacetate  
228 succinimidyl ester (CFDA/SE). OY-107 was stained  
229 using the red fluorescent stain 5-(and 6)-carboxyte-  
230 tramethylrhodamine succinimidyl ester (TAMRA/SE).  
231 Immediately prior to injection, cell suspensions were  
232 diluted into site groundwater to a nominal concen-  
233 tration of 1.5e8 cells/ml.

234 Bacterial cell counts were monitored using ferro-  
 235 graphic capture, which combines the selectivity of  
 236 immunomagnetic tagging with the high resolution of  
 237 ferrography. For the ferrographic capture analyses,  
 238 polyclonal rabbit antibodies (Rockland Immunochem-  
 239 icals, Gilbertsville, PA) raised to whole cells of the  
 240 target bacterial strains were used to tether goat anti-  
 241 rabbit-coated paramagnetic beads (50-nm diameter,  
 242 Miltenyi Biotec, Auburn, CA) to the surface of the  
 243 target cells following sample collection (Zhang and  
 244 Johnson, 1999; Zhang et al., 1999). The bacterium–  
 245 bead suspension was introduced into a Bio-Ferrograph  
 246 (Guilfoyle, Belmont, MA), which deposited the mag-  
 247 netically tagged bacteria onto a small area on a glass  
 248 substratum. The bacteria were then enumerated under  
 249 an epifluorescence microscope. TAMRA/SE- and  
 250 CFDA/SE-stained cells were enumerated using red  
 251 ( $\lambda_{\text{ex}} = 510\text{--}560$  nm, dichroic—575 nm,  $\lambda_{\text{em}} = 590$  nm)  
 252 and green ( $\lambda_{\text{ex}} = 470\text{--}490$  nm, dichroic—505 nm,  
 253  $\lambda_{\text{em}} = 510\text{--}530$  nm) filters, respectively. Previous  
 254 studies have shown that ferrographic capture provides  
 255 virtually 100% recovery of target cells and yields  
 256 extremely high analytical resolution ( $\sim 20$  cells/ml)  
 257 (Zhang and Johnson, 1999; Zhang et al., 1999, 2001;  
 258 Johnson et al., 2000, 2001).

259 The same two bacterial strains were used in the  
 260 year 2000 and year 2001 injections; however, the  
 261 stains were switched between the two strains for the  
 262 July 2001 experiment relative to the July 2000 experi-  
 263 ment (Table 1). Hence, in groundwater samples, the  
 264 two different strains were distinguished antigenically,  
 265 whereas the injection year was distinguished using the  
 266 internal stain. Unstained (or weakly stained) cells  
 267 were also monitored using anti-DA001 and anti-OY-  
 268 107 antibodies that had been conjugated to a green  
 269 fluorophore, fluorescein isothiocyanate (FITC).

270 Since the FITC-conjugated antibodies provide a  
 271 visible outline of both unstained and stained target  
 272 cells, the unstained cell concentrations were deter-  
 273 mined by difference between analyses using FITC-  
 274 conjugated antibodies (stained plus unstained cells)

275 and nonconjugated antibodies (stained cells). The  
 276 antibodies may have also captured antigenically sim-  
 277 ilar nontarget cells. Although nontarget cells would  
 278 not be enumerated using nonconjugated antibodies (in  
 279 analyses for stained cells), they would be enumerated  
 280 using FITC-conjugated antibodies (in analyses for  
 281 stained plus unstained cells). Hence, the “unstained”  
 282 cell counts may include both target strains as well as  
 283 antigenically similar cells. Laboratory tests showed  
 284 that the antibodies successfully capture wild-type  
 285 DA001 and wild-type OY-107 cells (where wild type  
 286 refers to initial isolates), indicating that the analyses  
 287 could indeed be capable of capturing indigenous  
 288 forms of these strains.

289 Since both bacterial strains were potentially present  
 290 in both colors in each groundwater sample, the extent  
 291 of antibody cross-reactivity between the two bacterial  
 292 strains was determined. Serial dilutions of both bacte-  
 293 rial strains collected from their respective injection  
 294 lines were analyzed using the respective nontarget  
 295 antibody. It was determined that approximately 1% of  
 296 cells from serial dilutions of both bacterial strains  
 297 reacted with their respective nontarget antibody (data  
 298 not shown). However, unstained cell suspensions  
 299 grown and serially diluted in the laboratory showed  
 300 zero cross-reactivity with their nontarget antibody  
 301 (data not shown), indicating that the apparent 1%  
 302 cross-reactivity observed in the injection line stand-  
 303 ards may have resulted from cross-contamination  
 304 during sampling of the injection lines. Even 1%  
 305 apparent cross-reactivity would have insignificant  
 306 effect on the bulk of our analyses, since cell concen-  
 307 trations examined by ferrographic capture were pri-  
 308 marily in the 10–1000 cells/ml range, yielding a  
 309 maximum of 10 cells/ml nontarget cells in the sam-  
 310 ples. However, interpretation of results from samples  
 311 containing greater than 1000 cells/ml necessarily  
 312 considered potential effects of cross-reactivity.

313 Another potential artifact, given the possibility that  
 314 each of the two strains could be present in each of the  
 315 two colors, was the potential visibility of one stain  
 316 under the filter used for the other stain during enu-  
 317 meration. To avoid this problem, filters were switched  
 318 back and forth during enumeration to ensure that the  
 319 cells being counted were exclusively observed under  
 320 the target filter.

321 Samples taken from the injection solutions were  
 322 collected to serve as standards (following dilution to

t1.1 Table 1  
 t1.2 Stains used during the series of experiments

t1.3	Year	DA001	OY-107
t1.4	2000	TAMRA/SE	CFDA/SE
t1.5	2001	CFDA/SE	TAMRA/SE

323 1000 cells/ml) for ferrographic analysis. These stand-  
 324 ards and blanks consisting of high purity Milli-Q  
 325 water were included in each set of eight samples  
 326 analyzed. Standard concentrations of both bacterial  
 327 strains were observed to degrade at approximately 1%  
 328 per day in all ferrographic analyses (using FITC-  
 329 conjugated and nonconjugated antibodies). The  
 330 decrease in captured cell number over time is likely  
 331 due to antigen alteration in the presence of form-  
 332 aldehyde, as has been previously reported (Johnson et  
 333 al., 2000).

334 Flow cytometry (FACScan, Becton-Dickinson  
 335 Immunocytometry Systems, San Jose, CA) was used  
 336 to monitor cell concentrations of CFDA/SE-stained  
 337 DA001 when they were present at concentrations  
 338 greater than 5000 cells/ml. Although ferrographic  
 339 capture can analyze cell concentrations above this  
 340 magnitude following 1:10 or greater dilution, flow  
 341 cytometry has the advantage of being extremely fast.  
 342 A filter to allow detection of TAMRA/SE-stained OY-  
 343 107 was not available on the instrument, and so  
 344 TAMRA/SE-stained cells were enumerated exclu-  
 345 sively using ferrographic capture. TransFluoSpheres  
 346 (Molecular Probes, Eugene, OR) were added at a  
 347 known concentration (60,000 spheres/ml) to the sam-  
 348 ples in order to determine volume of sample analyzed.  
 349 Each sample was run for 50 s at the low flow rate  
 350 setting of the machine, which corresponds to approx-  
 351 imately 10  $\mu$ l/min.

### 352 3. Field experiments

353 The South Oyster (SO) focus area is one of two  
 354 flow cells located at the South Oyster site in Oyster,  
 355 VA on the southern end of the Delmarva Peninsula.  
 356 The SO flow cell is located in a surficial aquifer that  
 357 comprised of unconsolidated to weakly cemented,  
 358 well-sorted, medium- to fine-grained sands and peb-  
 359 bly sands. The aquifer is underlain by a fine-grained  
 360 muddy silt aquitard situated approximately 9 m  
 361 below ground surface (BGS). A peat layer from  
 362 ~ 2.5 to 4.0 m BGS lies within the granular sands.  
 363 The SO focus area flow cell is bounded vertically by  
 364 the peat layer above and the clay aquitard below.  
 365 The sediment between the peat layer and the aquitard  
 366 is dominantly very fine sand and fine sand (quartz,  
 367 feldspar, and significant micaceous minerals) with

368 occasional medium- and coarse-grained sand string-  
 369 ers. The SO flow cell is characterized as suboxic,  
 370 with dissolved oxygen concentrations in the range of  
 371 0.2–1.0 mg/l.

372 Twenty-four multilevel samplers (MLS) exist in  
 373 the flow cell (Fig. 1) in an array that is oriented with  
 374 the main axis parallel to natural local groundwater  
 375 flow direction, as inferred from water level observa-  
 376 tions prior to site installation. The flow cell is  
 377 bounded at the up-gradient limit by the injection well  
 378 SO-B2 and at the down-gradient limits by three  
 379 extraction wells SO-A3, SO-B3, and SO-C3, which  
 380 form a line perpendicular to flow about 19.5 m down-  
 381 gradient of SO-B2.

382 Each MLS consisted of a 3-cm-diameter PVC rod  
 383 separated into 12 equally spaced zones that is sepa-  
 384 rated by neoprene baffles. Stiff polyethylene tubing  
 385 (3.2 mm ID) extended from approximately 0.5 m  
 386 above ground surface to each sampling port. Detailed  
 387 descriptions of the MLS used at SO focus area can be  
 388 found elsewhere (Mailloux et al., in press). In this  
 389 paper, the MLS are denoted by their number and the  
 390 prefix “SO”, e.g. SO-17 denotes South Oyster focus  
 391 area MLS 17. The sampling port is denoted by




Fig. 1. Layout of the South Oyster focus area flow cell. Numbers refer to multilevel samplers (MLS). B2 is the injection MLS. The forced gradient was set 1 week prior to injection by extraction from three wells in a line oriented perpendicular to flow 12 m down-gradient of MLS SO-24.

392 number (1–12) and the prefix “P”, e.g. SO-17 P4  
393 denotes port 4 in MLS 17. The port number increases  
394 with depth in the flow cell.

395 The ambient pore water velocity at the site is  
396 approximately 0.1 m/day. Forced gradient conditions  
397 were established at the site approximately 48 h prior  
398 to the start of injection by withdrawing from the three  
399 down-gradient extraction wells, SO-A3, SO-B3, and  
400 SO-C3, at flow rates of 20, 40, and 20 ml/min,  
401 respectively. The resulting average pore water veloc-  
402 ity was approximately 1 m/day.

403 Injection was conducted in MLS SO-B2, a 4-in.  
404 PVC well with four vertically discrete injection  
405 zones separated by packers. In contrast to the  
406 previous experiment (July 2000), in which both  
407 bacterial strains were injected over the entire depth  
408 of the flow cell, the July 2001 experiment injected  
409 the two bacterial strains over two discreet depths.  
410 DA001 was injected over the interval of  $-3.5$  to  
411  $-4.5$  m relative to mean sea level (MSL), and OY-  
412 107 was injected over the interval of  $-5.0$  to  
413  $-6.0$  m MSL, both at injection rate of 0.8 l/min.  
414 The bacterial injection zones were separated by  
415 injection of unamended groundwater into the inter-  
416 val from  $-4.5$  to  $-5.0$  m MSL at a rate of 0.4 l/  
417 min. Unamended groundwater was also injected  
418 above the DA001 injection zone ( $-3.0$  to  $-3.5$   
419 m MSL) at a rate of 0.4 l/min. The total injection  
420 rate from SO-B2 was 2.5 l/min. Injection occurred  
421 over a 12-h period. Both bacterial injection solu-  
422 tions were sampled every 30 min during the injec-  
423 tion interval.

424 A custom manifold was developed to apply con-  
425 tinuous suction to all sampled ports simultaneously  
426 using a peristaltic pump (Johnson et al., 2000; Mail-  
427 loux et al., in press). All sampled ports were pumped  
428 continuously and equally in order to minimize induce-  
429 ment of a vertical hydraulic gradient between the  
430 ports. Relative to the previous experiment (July  
431 2000), the number of samples taken during early  
432 breakthrough was increased in the July 2001 experi-  
433 ment to increase the resolution of the early break-  
434 through pulses. All bacterial samples were taken using  
435 50-ml polypropylene centrifuge tubes and then fixed  
436 with 1% formaldehyde. Samples were then split into  
437 smaller aliquots (15 ml), briefly stored on ice, shipped  
438 back to the University of Utah on ice (unfrozen), and  
439 stored at 4 °C.

#### 4. Laboratory experiments

440

441 Several laboratory experiments were run to inves-  
442 tigate potential enhanced detachment of attached cells  
443 in a controlled system. Each experiment involved a  
444 loading step, in which a bacterial suspension was  
445 introduced to a sediment-packed column. Loading  
446 was followed by elution until the effluent cell con-  
447 centration was 100 cells/ml or less. Elution was  
448 followed by injection of a cell suspension to deter-  
449 mine whether detachment of previously loaded (resi-  
450 dent) cells was enhanced by arrival of mobile cells.

451 Sediments used in the column experiments were  
452 obtained from the narrow channel (NC) focus area in  
453 Oyster, VA. The sediment is primarily composed of  
454 quartz, feldspar, clays, and iron and aluminum  
455 hydroxides in decreasing order of abundance (John-  
456 son et al., 2000). A target average pore water velocity  
457 of 1 m/day for each experiment was maintained  
458 throughout the duration of the experiment. Flow was  
459 allowed to equilibrate at the target average pore water  
460 velocity for at least 1 pore volume (PV) prior to  
461 injection of the bacterial solutions. Narrow channel  
462 artificial groundwater (NCAGW) was used as the  
463 aqueous medium in all laboratory transport experi-  
464 ments (DeFlaun et al., 2001).

465 Strains were grown according to growth protocols  
466 described earlier (DeFlaun et al., 2001). Following  
467 collection, all samples were fixed with 1% formalde-  
468 hyde. Since the concentrations of attached resident  
469 cells in the field experiments were not known, it was  
470 not clear what attached bacterial concentrations would  
471 be reflective of field conditions. For this reason,  
472 injection duration (0.2–0.5 pore volumes), injection  
473 concentration ( $2 \times 10^3$  to  $4 \times 10^5$  cells/ml), and column size  
474 (2.5 cm diameter by 15 cm length to 7 cm diameter by  
475 53 cm length) were varied among the set of column  
476 experiments to yield resident (attached) cell concen-  
477 trations ranging from about 50 to about  $1.5 \times 10^5$  cells per  
478 gram of sediment. The concentration of subsequently  
479 injected bacteria was also varied ( $2 \times 10^3$  to  $4 \times 10^5$  cells/ml).

#### 5. Results

480

481 MLS SO-10 and SO-14 were sampled in July  
482 2001, prior to the establishment of forced gradient  
483 conditions at the site, in order to determine the stained  
484

484 resident cell concentrations (remaining from July  
485 2000 injection). Stained resident cells of both strains  
486 remaining from the July 2000 injection were observed  
487 in both sampled MLS. TAMRA/SE-stained DA001  
488 concentrations were approximately 50 cells/ml at all  
489 depths in SO-10, and were approximately 200 cells/ml  
490 at all depths in SO-14. CFDA/SE-stained resident OY-  
491 107 cells were observed at low concentrations (aver-  
492 aging ~ 20 cells/ml) at all depths in SO-10 and low  
493 concentrations (averaging ~ 30 cells/ml) at all depths  
494 in SO-14. Laboratory blanks showed no cells, proving  
495 that the cells did not derive from contamination during  
496 analysis. Tubing in all MLS was removed and  
497 replaced prior to monitoring for the year 2001 injection.  
498 eliminating the possibility that these cells had  
499 remained in the tubing during the year that passed  
500 between experiments.

501 No enhanced detachment of stained resident cells  
502 was observed to occur with the arrival of injected cells  
503 during the July 2001 injection. This is illustrated in  
504 several sampling ports in various MLS, including SO-  
505 T2 P12, SO-17 P10, SO-24 P8, and SO-24 P12 (Fig.

2), where breakthrough of injected DA001 at 70 (SO-  
T2 P12), 35 (SO-17 P10), and 160 (SO-24 P8 and  
P12) elapsed hours was not accompanied by increases  
in stained resident cell concentrations. SO-T2 P12  
shows an example of the background concentrations  
of stained resident cells in samples taken prior to  
breakthrough of injected cells. The background con-  
centrations of stained resident cells were not observed  
in all of the MLS, including three of the four MLS  
shown in Fig. 2. Results from four other monitored  
MLS (not shown) also showed a lack of enhanced  
detachment of stained resident cells regardless of  
presence of stained resident cells, indicating that  
enhanced detachment of stained resident cells did  
not occur in response to breakthrough of injected  
cells.

It should be noted that the analytical resolution of  
the analysis allowed enumeration of extremely low  
cell concentrations during initial breakthrough, and  
that variations among concentrations during initial  
breakthrough represent a combination of transport  
effects due to aquifer heterogeneity, sampling errors,




Fig. 2. Results from July 2001 field experiment showing breakthrough of injected DA001 and OY-107. Detachment of resident DA001 and OY-107 cells was not enhanced by breakthrough of the injected cells. Resident DA001 was observed in the aqueous phase before, during, and after extraction (e.g. SO-T2 P12).

528 and analytical error. In addition to high resolution, the  
 529 method provided sufficient flexibility to allow simul-  
 530 taneous monitoring of two strains each of two differ-  
 531 ent colors (each of two different injection times) as  
 532 well as their unstained counterparts in combination  
 533 with any antigenically similar strains.

534 Unstained cell concentrations showed ephemeral  
 535 increases coincident with the arrival of stained in-  
 536 jected cells, similarly to increases previously reported  
 537 from the July 2000 injection (Johnson et al., 2001).  
 538 MLS SO-17 P2 (Fig. 3) shows, in log (left) and  
 539 linear (right) format, ephemeral increases of total  
 540 (injected plus unstained) cells. These ephemeral  
 541 increases were a factor of three to four times greater  
 542 than injected cell concentrations in pulses centered at  
 543 70, 98, and 120 elapsed hours. MLS SO-17 P4 (Fig.  
 544 3) shows ephemeral increases of total cells, a factor  
 545 of three to four times greater than injected cell  
 546 concentrations in pulses centered at 106 and 145  
 547 elapsed hours.

548 The unstained cells were observed only in  
 549 limited locations throughout the site and were

550 exclusively observed in relatively shallow MLS  
 551 ports (port 4 and above). No significant increases  
 552 in total relative to injected cell concentrations were  
 553 observed in the majority of ports analyzed, as  
 554 shown in MLS SO-17 P6, SO-24 P8, and SO-24  
 555 P10 (Fig. 4), where stained and total cell concen-  
 556 trations increased simultaneously with negligible  
 557 differences (less than a factor of two) upon arrival  
 558 of injected cells.

559 Results from the laboratory experiments also  
 560 indicate that no enhanced detachment of resident  
 561 cells occurred in response to the arrival of injected  
 562 cells. Results from a representative experiment are  
 563 shown in Fig. 5. One pore volume of stained DA001  
 564 ( $2e5$  cells/ml) was injected into the column (0–2  
 565 elapsed hours), resulting in an effluent pulse from 2  
 566 to 5 elapsed hours. Extended tailing of low concen-  
 567 trations of stained DA001 was observed during  
 568 elution with NCAGW. In this column experiment,  
 569 two separate reservoirs were used for delivery of the  
 570 bacterial suspension and NCAGW, and switching  
 571 between the two reservoirs required the use of a




Fig. 3. Results from July 2001 field experiment showing an ephemeral pulse of unstained cells (difference between total and stained cell concentrations) coincident with the breakthrough of injected cells. The peak unstained cell concentrations were more than a factor of two greater than the concentrations of injected cells in several ports. This result was observed only in the shallow ports.




Fig. 4. Results from the July 2001 field experiment highlighting the lack of significant unstained cell concentrations (difference between total and stained cells) in the majority of the MLS ports sampled at the site.

572 three-way valve. Unlike the field experiment, switching  
 573 between reservoirs during the laboratory experiment  
 574 may have caused a sudden increase in hydrodynamic  
 575 shear in the column. To assess the magnitude of  
 576 enhanced detachment due to hydrodynamic shear,  
 577 the three-way valve was quickly switched between  
 578 reservoirs at about 24 elapsed hours (Fig. 5).  
 579 hours (Fig. 5).

580 Following the valve switch, a pulse injection of  
 581  $3.5 \times 10^5$  cells/ml of unstained DA001 cells was  
 582 then introduced into the column beginning at  $\sim 29$   
 583 elapsed hours. The effluent pulse of stained resident  
 584 cells at 24–28 elapsed hours ( $\sim 1 \times 10^3$  cells/ml) was

585 associated with the valve switch. The effluent pulse  
 586 of stained resident cells at 30–33 elapsed hours ( $1 \times 10^2$   
 587 cells/ml) was associated with the breakthrough of  
 588 unstained DA001 cells. The fact that greater  
 589 enhanced detachment of resident cells occurred  
 590 during the valve switch alone relative to the valve  
 591 switch plus introduction of mobile cells indicates  
 592 that the observed enhanced detachment of the resi-  
 593 dent cells in the laboratory experiment may have  
 594 resulted from hydrodynamic shear associated with  
 595 the valve switch, and cannot be attributed to colli-  
 596 sion with mobile cells. It should be noted that  
 597 subsequent experiments conducted in a system that




Fig. 5. Representative results from the laboratory column experiments showing a lack of enhanced detachment of stained resident cells. The first two vertical bars represent the time of the injection pulse for stained DA001. The third vertical bar represents the time of the valve switch. The fourth and fifth vertical bars represent the time of the injection pulse of unstained DA001 cells.

598 avoided valve switching also lacked enhanced de-  
599 tachment of resident cells.

## 600 6. Discussion

601 Enhanced detachment of stained resident cells in  
602 response to arrival of injected cells was not observed  
603 in the July 2001 field injection, nor was it observed in  
604 any of the laboratory experiments. Lack of enhanced  
605 detachment of stained resident cells may indicate that  
606 the concentrations of stained resident (attached) cells  
607 were insufficient to support significant enhanced  
608 detachment in the presence of mobile cells, assuming  
609 that greater attached cell concentrations increase the  
610 magnitude of enhanced detachment. The adhesion-  
611 deficient variants used in the experiments may have  
612 yielded low attached cell concentrations relative to  
613 their indigenous counterparts, although the attached  
614 concentrations of the latter are unknown.

615 Alternatively, the results may indicate that the  
616 ephemeral pulses of unstained cells observed during  
617 the field experiments represent a process other than  
618 enhanced detachment. Mechanisms, other than  
619 detachment, that may explain the ephemeral increases  
620 in unstained cell concentrations (coincident with the  
621 initial breakthrough of injected cells) include prefer-  
622 ential stain loss from the injected cell population  
623 residing in the low-concentration fringe of the plume

(Johnson et al., 2001). Stain loss by diffusion from  
624 cells could occur preferentially from cells residing at  
625 the low-concentration fringe of the injected plume.  
626 However, Fig. 6 shows that CFDA/SE-stained DA001  
627 monitored using FITC-conjugated antibodies (total  
628 cells) and nonconjugated antibodies (stained cells)  
629 degraded at very similar rates, indicating that stain  
630 loss over time was negligible.

631  
632 Notably, injection-zone samples taken during  
633 injection showed concentrations of  $1.3 \times 10^8$  and  $1.2 \times 10^8$   
634 cells/ml for DA001 and OY-107, respectively, ac-  
635 cording to ferrographic capture using FITC-conju-  
636 gated antibodies. This result agreed well with flow  
637 cytometry results for samples collected from the  
638 injection tanks, which showed DA001 concentrations  
639 of  $1.5 \times 10^8$  cells/ml. Surprisingly, concentrations of  
640 both strains determined ferrographically using non-  
641 conjugated antibodies were about  $\sim 55\%$  of those  
642 determined by the other methods. This latter result  
643 was corroborated by direct counts on filters (Mark  
644 Fuller, Envirogen, personal communication). These  
645 results indicate that nearly 50% of the injected cells  
646 were weakly stained. The weakly stained cells in the  
647 injection solution may represent inefficiency in the  
648 staining procedure (Mark Fuller, Envirogen, personal  
649 communication).

650 The presence of unstained cells in the injected  
651 solution does not affect the analysis presented  
652 regarding the detachment of stained resident cells.  
653 The presence of unstained cells in the injection




Fig. 6. CFDA/SE-stained DA001 standard results observed using FITC-conjugated antibodies (total DA001) and nonconjugated antibodies (stained DA001) to DA001 cells.

654 solution does, however, explain the persistent factor-  
655 of-two difference between total and unstained cell  
656 concentrations observed in nearly all ports when cell  
657 concentrations became significant (e.g. greater than  
658 100 cells/ml in Figs. 3 and 4), as well as the factor-  
659 of-two difference between standards monitored using  
660 FITC-conjugated versus nonconjugated antibodies  
661 (Fig. 6). The presence of unstained cells in the  
662 injection solution does not, however, explain the  
663 ephemeral pulses of unstained cells yielding ratios  
664 of total to injected cell concentrations of three to  
665 four (Fig. 3).

666 Stain loss due to cell division at the low-concen-  
667 tration fringe of the plume of injected bacteria repre-  
668 sents another potential means of generating the  
669 ephemeral pulses of unstained cells. Recall that the  
670 ephemeral pulses of unstained cells were observed in  
671 relatively shallow ports. Indeed, cells in samples  
672 collected from shallow ports exhibited a polar distri-  
673 bution of stain that was suggestive of stain loss due to  
674 cell division, in which the daughter cells each retained  
675 stain in the portion of the cell that was originally part  
676 of the parent cell. Fig. 7 shows two images of polar  
677 stained cells (left) and normal fully stained cells  
678 (right). The cells that exhibited a polar distribution  
679 of stain would easily be missed during enumeration  
680 following ferrographic capture by nonconjugated anti-  
681 bodies due to their relatively weak illumination. It is  
682 therefore possible that the ephemeral pulses of  
683 unstained cells coincident with the injected cells  
684 represent stain loss due to cell division in the low-  
685 concentration fringe of the plume in the shallow  
686 portion of the flow cell.

687 Cell division in the low-concentration fringe of the  
688 plume could potentially have been driven by a favor-  
689 able combination of electron acceptor and donor  
690 exclusively at the interface of the injected and ambient  
691 groundwater in the shallow portion of the aquifer.  
692 Differences in the chemistries of the injected versus  
693 ambient groundwater, if any, are expected to have  
694 been subtle, since the injected groundwater was  
695 extracted from the site at a location just outside the  
696 flow cell. The limited groundwater chemistry data  
697 measured during the field injection indicate no sig-  
698 nificant variation with depth for the measured param-  
699 eters (including dissolved oxygen, dissolved organic  
700 carbon, ionic strength, and pH). The limited data do  
701 suggest, however, that dissolved ( $<0.2 \mu\text{m}$ ) organic  
702 carbon concentrations were slightly higher in the  
703 ambient relative to the injected groundwater (by 1–  
704 10 mg/l), potentially supporting growth at the inter-  
705 face. However, the temporal and spatial sparseness of  
706 the chemical data prevent rigorous identification of  
707 potential chemical drivers for cell division at the  
708 interface between ambient and injected groundwater,  
709 and so this determination is beyond the capabilities of  
710 the present study.

711 This investigation highlights the resolution of fer-  
712 rographic capture, which clearly showed that  
713 enhanced detachment of stained resident cells did  
714 not occur in response to breakthrough of injected cells  
715 in our system. This result weakens the original  
716 hypothesis that hydrodynamic collision between  
717 mobile and attached bacteria resulted in the observed  
718 ephemeral domination by unstained cells during initial  
719 breakthrough.




Fig. 7. Images of DA001 cells from the July 2001 field experiment showing polar distribution of stain in relatively dim cells, suggestive of cell division (a). Staining normally results in homogeneously bright cells (b).

720 **7. Uncited reference**

721 Hendry et al., 1999

722 **Acknowledgements**

723 This investigation was supported by the US  
 724 Department of Energy (DOE), Natural and Accelerated  
 725 Bioremediation Research Program (NABIR)-  
 726 Acceleration Element (Grant DE-FG03-99ER62820/  
 727 A000). The authors greatly appreciate the leadership  
 728 of Dr. Frank Wobber, the Program Manager for the  
 729 Acceleration Element of NABIR. Access of the field  
 730 site was granted by the Nature Conservancy, Virginia  
 731 Coast Reserve. We also acknowledge the excellent  
 732 work of University of Utah undergraduate students  
 733 Tyler Knudsen, Suzanne Bethers, David Stone, Brad  
 734 Murray, and the rest of the team FASTBUGS  
 735 (Ferrographic Analysis in Short Time by Under-  
 736 graduate Students). Finally, we wish to acknowledge  
 737 the superb efforts of Mark Fuller, Tim Scheibe, Brian  
 738 Mailloux, Tim Griffin, Mary DeFlaun, and T.C.  
 739 Onstott in the planning and execution of the field  
 740 work performed at the South Oyster site.

741 **References**

742 Bergendahl, J., Grasso, D., 2000. Prediction of colloid detachment  
 743 in a model porous media: hydrodynamics. *Chemical Engineer-*  
 744 *ing Science* 55, 1523–1532.  
 745 Dabros, T., 1989. Interparticle hydrodynamic interactions in depo-  
 746 sition processes. *Colloids and Surfaces. A, Physicochemical and*  
 747 *Engineering Aspects* 39, 127–141.  
 748 Dabros, T., van de Ven, T.G.M., 1992. Hydrodynamic interactions  
 749 between two spheres near a solid plane. *International Journal of*  
 750 *Multiphase Flow* 18, 751–764.  
 751 DeFlaun, M.F., Tanzer, A., McAteer, A., Marshall, B., Levy, S.,  
 752 1990. Development of an adhesion assay and characterization of  
 753 an adhesion-deficient mutant of *Pseudomonas fluorescens*. *Ap-*  
 754 *plied and Environmental Microbiology* 56, 112–119.  
 755 DeFlaun, M.F., Fuller, M.E., Zhang, P., Johnson, W.P., Mailloux,  
 756 B.J., Holben, W.E., Kovacic, W.P., Balkwill, D.L., Onstott,  
 757 T.C., 2001. Comparison of innovative methods for monitoring  
 758 bacterial transport. *Journal of Microbiological Methods* 47,  
 759 219–231.  
 760 Fuller, M.E., Mailloux, B.J., Zhang, P., Streger, S.H., Hall, J.A.,  
 761 Vainberg, S.N., Beavis, A.J., Johnson, W.P., Onstott, T.C.,  
 762 DeFlaun, M.F., 2001a. Evaluation of CFDA/SE-staining  
 763 coupled with multiple detection methods during a field-scale

bacterial transport experiment. *FEMS Microbiology, Ecology*  
 37, 55–66. 764  
 765  
 Fuller, M.E., Streger, S.H., Rothmel, R.K., Mailloux, B.J., Hall,  
 J.A., Onstott, T.C., Fredrickson, J.K., Balkwill, D.L., DeFlaun,  
 M.F., 2001b. Development of a vital fluorescent staining  
 method for monitoring bacterial transport in subsurface en-  
 vironments. *Applied and Environmental Microbiology* 66,  
 4486–4496. 766  
 767  
 768  
 769  
 770  
 771  
 Harvey, R.W., 1997. Microorganisms as tracers in groundwater  
 injection and recovery experiments: a review. *FEMS Microbi-*  
 ology Reviews 20, 461–472. 772  
 773  
 774  
 Harvey, R.W., Garabedian, S.P., 1991. Use of colloid filtration  
 theory in modeling movement of bacteria through a contami-  
 nated sandy aquifer. *Environmental Science and Technology* 25,  
 178–185. 775  
 776  
 777  
 778  
 Hendry, M.J., Lawrence, J.R., Maloszewski, P., 1999. Effects of  
 velocity on the transport of two bacteria through saturated sand.  
*Ground Water* 37, 103–112. 779  
 780  
 781  
 Johnson, W.P., Blue, K.A., Logan, B.E., Arnold, R.G., 1995. Mod-  
 eling bacterial detachment during transport through porous me-  
 dia as a residence-time-dependent process. *Water Resources*  
*Research* 31, 2649–2658. 782  
 783  
 784  
 785  
 Johnson, W.P., Zhang, P., Fuller, M.E., Scheibe, T.D., Mailloux,  
 B.J., Onstott, T.C., DeFlaun, M.F., Hubbard, S.S., Radtke, J.,  
 Kovacic, W.P., Holben, W., 2000. Ferrographic tracking of  
 bacterial transport in the field at the narrow channel focus  
 area, Oyster, VA. *Environmental Science and Technology* 35,  
 182–191. 786  
 787  
 788  
 789  
 790  
 791  
 Johnson, W.P., Zhang, P., Gardner, P.M., Fuller, M.E., DeFlaun,  
 M.F., 2001. Evidence for detachment of indigenous bacteria  
 from aquifer sediment in response to arrival of injected bacteria.  
*Applied and Environmental Microbiology* 67, 4908–4913. 792  
 793  
 794  
 795  
 Mailloux, B.J., Fuller, M.E., Rose, G.F., Onstott, T.C., DeFlaun,  
 M.F., Alvarez, E., Hemingway, C., Hallet, R.B., Phelps, T.J.,  
 Griffin, T., 2002. Design of an integrated injection and sam-  
 pling system for groundwater tracer experiments. *Ground*  
*Water* (in press). 796  
 797  
 798  
 799  
 800  
 Meinders, J.M., van der Mei, H.C., Busscher, H.J., 1995. Deposi-  
 tion efficiency and reversibility of bacterial adhesion under flow.  
*Journal of Colloid and Interface Science* 176, 329–341. 801  
 802  
 803  
 Ryan, J.N., Elimelech, M., 1996. Colloid mobilization and transport  
 in groundwater. *Colloids and Surfaces. A, Physicochemical and*  
*Engineering Aspects* 107, 1–56. 804  
 805  
 806  
 Steffan, R.J., Sperry, K.L., Walsh, M.T., Vainberg, S., Condee,  
 C.W., 1999. Field-scale evaluation of in situ bioaugmentation  
 for remediation of chlorinated solvents in groundwater. *Environ-*  
*mental Science and Technology* 33, 2771–2781. 807  
 808  
 809  
 810  
 Zhang, P., Johnson, W.P., 1999. Rapid selective ferrographic enu-  
 meration of bacteria. *Journal of Magnetism and Magnetic Ma-*  
*terials* 194, 267–274. 811  
 812  
 813  
 Zhang, P., Johnson, W.P., Rowland, R., 1999. Bacterial tracking  
 using ferrographic separation. *Environmental Science and Tech-*  
*nology* 33, 2456–2460. 814  
 815  
 816  
 Zhang, P., Johnson, W.P., Scheibe, T.D., Choi, K.H., Dobbs, F.C.,  
 Mailloux, B.J., 2001. Extended tailing of bacteria following  
 breakthrough at the narrow channel focus area, Oyster, VA.  
*Water Resources Research* 37, 2687–2698. 817  
 818  
 819  
 820